









KEMH ROTEM Algorithm for Critical Bleeding

Key Points: This algorithm should be used in conjunction with the KEMH Blood Product Guidelines for Major Obstetric Haemorrhage. Only treat abnormal values if active bleeding or at high risk of bleeding. Consider early tranexamic acid (1g IV) in all critical bleeding situations. Repeat ROTEM analysis 10 mins after intervention to assess response.

	ABNORMAL ROTEM	CRITERIA	DIAGNOSIS	INTERVENTION	CORRECTED ROTEM
FIBRINOGEN		FIBTEM A5 ≤ 10 mm	Low fibrinogen	Cryoprecipitate OR fibrinogen concentrate (see dosing guide) AND tranexamic acid 1 g (if not already administered)	
PLATELETS		EXTEM A5 ≤ 35 mm and FIBTEM A5 ≥ 10 mm	Low platelets	Platelets: 1 adult dose (correlate with platelet count)	
		EXTEM A5 ≤ 25 mm and FIBTEM A5 ≤ 10 mm	Low platelets and Low fibrinogen	Platelets and fibrinogen (correlate with platelet count)	
FACTORS		EXTEM CT 80-140s and FIBTEM A5 ≤ 10 mm	Low fibrinogen	Correct fibrinogen and reassess	
		EXTEM CT >140 s and FIBTEM A5 ≤ 10 mm	Low fibrinogen and Low coagulation factors	FFP 1-2U + fibrinogen as indicated (Consider Prothrombinex-see below)	
FIBRINOLYSIS		Early Diagnosis EXTEM A5 ≤ 35 mm or FIBTEM CT >80 s	High likelihood of excess fibrinolysis	Tranexamic acid 1g Consider repeat dose if has lost over 1 blood volume since initial dose	
		Late Diagnosis EXTEM or FIBTEM ML $\geq 5\%$	Excess fibrinolysis		

Fibrinogen Dosing Guide

FIBTEM A5 Target: ≥ 12 mm

FIBTEM A5	Cryoprecipitate	Fibrinogen Concentrate
7-10mm	1-2 doses	2g
≤ 6 mm	2 doses	4g

Cryoprecipitate is the default product for fibrinogen replacement in the majority of bleeding cases at KEMH. It is generally available within 10 minutes of the request being made. One dose is equivalent to 10 whole blood units or 5 apheresis units and may be supplied as a combination of these units.

Fibrinogen Concentrate

Guidelines For Use

- Should only be considered in exceptional circumstances and with consultant anaesthetist or haematologist approval.
- Fibrinogen concentrate may be indicated instead of, or in addition to, cryoprecipitate if the FIBTEM A5 is ≤ 6 mm in patients experiencing life threatening bleeding, or if there is critical bleeding with a high suspicion of coagulopathy.
- Use at higher FIBTEM values may be appropriate in patients refusing cryoprecipitate.

Administration

- Reconstitute 1g in 50ml warm sterile water (use prepared kit in fluid warmer). Swirl gently and do not shake (to avoid foaming). Administer each 1g via syringe driver over 2-4 mins if life-threatening haemorrhage or over 10 mins if not.

Prothrombinex









- Haematologist approval required
- Consider as an alternative to FFP for patients with coagulation factor deficiency (e.g. prolonged EXTEM CT see above) in the following circumstances:
 - Circulatory overload
 - Rapid correction in extreme coagulopathy

Key Contact Information

- Bloodbank: dial 82748
- Duty Anaesthetist: *41225 (internal) or 0420 302 571
- Theatre co-ordinator: *41220 (internal) or 0424 148 574
- On call haematologist: via switchboard on 08 6458 2222
- Theatres 1-4: 82211, 82212, 82213 and 82214 respectively. Theatre 5: dial 81482
- ASCU dial 82155/6, Delivery suite co-ordinator: pager 3313

SCGH ROTEM Algorithm for Critical Bleeding

Key Points: This algorithm should be used in conjunction with the SCGH Critical Bleeding Protocol. Only treat abnormal values if active bleeding or at high risk of bleeding. Repeat ROTEM analysis 10 mins after intervention to assess response.


	ABNORMAL ROTEM	CRITERIA	DIAGNOSIS	INTERVENTION	CORRECTED ROTEM
FIBRINOLYSIS		Early Diagnosis EXTEM A5 ≤ 35mm or FIBTEM CT > 600s	High likelihood of excess fibrinolysis	Tranexamic acid 1g Consider repeat dose if has lost over 1 blood volume since initial dose	
		Late Diagnosis EXTEM or FIBTEM ML ≥ 5%	Excess fibrinolysis		
FIBRINOGEN		FIBTEM A5 ≤ 10mm	Low fibrinogen	Cryoprecipitate (see dosing guide)	
PLATELETS		EXTEM A5 ≤ 35mm and FIBTEM A5 > 10mm	Low platelets	Platelets: 1 adult dose (correlate with platelet count)	
		EXTEM A5 ≤ 25mm and FIBTEM A5 ≤ 10mm	Low platelets and Low fibrinogen	Platelets and fibrinogen (correlate with platelet count)	
FACTORS		EXTEM CT 80-140s and FIBTEM A5 ≤ 10mm	Low fibrinogen	Correct fibrinogen and reassess	
		EXTEM CT > 80s but FIBTEM A5 > 10mm	Low coagulation factors	FFP 1-4U or Prothrombinex 10 U/kg (+ fibrinogen if indicated)	
		EXTEM CT > 140s and FIBTEM A5 ≤ 10mm	Low fibrinogen and Low coagulation factors		

Fibrinogen Dosing Guide

FIBTEM A5 Target: ≥12mm

FIBTEM A5	Increase required	Cryoprecipitate*
9-10mm	2-3 mm	10 Units
7-8mm	4-5 mm	15 Units
4-6mm	6-8 mm	20 Units
<4mm	≥9mm	20-25 Units

*Cryoprecipitate dosing is for standard adult units
(Cryo 5 units = Fibrinogen increase of approx 2mm)











Prothrombinex

- Warfarin Reversal: Indicated for urgent reversal of warfarin in critical bleeding, usual dose 25-50U/kg (+/- FFP) discuss with haematologist.
- Consider as an alternative to FFP for patients with coagulation factor deficiency (e.g. prolonged EXTEM CT see above) in the following circumstances:
 - Circulatory overload
 - Rapid correction in extreme coagulopathy
 - Consider lower dose 10U/kg (round to nearest 500U).

FSH ROTEM Algorithm for Critical Bleeding

This algorithm should be used in conjunction with the FSH Major Haemorrhage Protocol
 Treat abnormal values only if there is active bleeding or the patients is at high risk of bleeding.
 Repeat ROTEM analysis 10 mins after any intervention to assess response.


	ABNORMAL ROTEM	CRITERIA	DIAGNOSIS	INTERVENTION	CORRECTED ROTEM
FIBRINOLYSIS		Trauma (within 3hrs) OR Post partum haemorrhage Flat trace OR Maximal lysis >5%	→ Hyperfibrinolysis	Tranexamic acid 1g	
FIBRINOGEN		FIBTEM A5 ≤10mm	Hypofibrinogenaemia	Cryoprecipitate	
PLATELETS		EXTEM A5 ≤35mm with normal fibrinogen*	Thrombocytopenia	Platelets	
FACTORS		EXTEM CT 90-140sec with normal fibrinogen** OR EXTEM CT >140sec	Low coagulation factors	Fresh Frozen Plasma 2-4u OR Prothrombinex 25IU/kg	

Cryoprecipitate Dosing Guide

FIBTEM A5	Non-obstetric	Obstetric
7-10	1 dose	2 doses
<6	2 doses	3 doses


One dose = five apheresis units = Fibrinogen A5 increase of approximately 4mm

*If EXTEM ≤25 and FIBTEM A5 ≤10 consider replacing both factors
 **Fibrinogen replacement in the context of hypofibrinogenaemia may overcome a minor prolongation of clotting time



Key components

EXTEM CT Clotting Time	Thrombin generation
EXTEM A5 Amplitude at 5 minutes	Fibrinogen and platelet concentration and function
FIBTEM A5 Amplitude at 5 minutes	Fibrinogen concentration and function
ML % Maximal lysis	Degree of fibrinolysis over temogram



Please stick this label in the patients progress notes

ROTEM ANALYSIS AND TREATMENT PLAN

Nurse or JMO to circle algorithm used then insert results from ROTEM Next circle range (action red range) and use algorithm to create a plan.

Date: / / Time:

ALGORITHM USED (circle one):
CARDIAC/VASCULAR or GENERAL/OBSTETRIC

- For CARDIAC/VASCULAR start here and do all:
INTEM CT = Below 205 / 205 & Above
HEPTEM CT = Below 205 / 205 & Above

- For GENERAL/OBSTETRIC start here (this section only):
EXTEM A5 = Below 35 / 35-40 / Above 40
FIBTEM A5 = Below 10 / 10-15 / Above 15
EXTEM CT = Below 80 / 80-90 / Above 90
EXTEM ML = Below 5 / 5 & Above

Management Plan:

Please stick this label in the patients progress notes